MUCINEX-SINUS MAX SEVERE CONGESTION AND PAIN CLEAR AND COOL AND MUCINEX NIGHTSHIFT SINUS CLEAR AND COOL- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, triprolidine hydrochloride RB Health (US) LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Mucinex Sinus-Max Severe Congestion & Pain Clear & Cool and Mucinex Nightshift Sinus Clear & Cool

Drug Facts

Uses

Mucinex Sinus-Max SEVERE CONGESTION & PAIN CLEAR & COOL temporarily relieves:

- nasal congestion
- headache
- minor aches and pains
- sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Mucinex Nightshift Sinus Clear & Cool temporarily relieves these common cold and flu symptoms:

- cough
- nasal congestion
- minor aches and pains

- sore throat
- headache
- sinus congestion and pressure
- runny nose
- ■sneezing
- itching of the nose or throat
- itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning (Nightshift Sinus Clear & Cool only):If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease heart disease diabetes
- high blood pressure
- thyroid disease
- glaucoma (Nightshift Sinus Clear & Cool only)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis (Nightshift Sinus Clear & Cool only)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (Nightshift Sinus Clear & Cool only)

When using this product **■** do not use more than directed

- excitability may occur, especially in children (Nightshift Sinus Clear & Cool only)
- marked drowsiness may occur (Nightshift Sinus Clear & Cool only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (Nightshift Sinus Clear & Cool only)
- avoid alcoholic drinks (Nightshift Sinus Clear & Cool only)
- use caution when driving a motor vehicle or operating machinery (Nightshift Sinus Clear & Cool only)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

MUCINEX SINUS-MAX SEVERE CONGESTION & PAIN CLEAR & COOL

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours

MUCINEX NIGHTSHIFT SINUS CLEAR & COOL

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use
- each 20 mL contains: sodium 9mg (Severe Congestion & Pain Clear & Cool only) and sodium 16mg (Nightshift Sinus Clear & Cool only)
- store at 20-25°C (68-77°F)
- do not refrigerate

(Mucinex Sinus-Max Severe Congestion & Pain Clear & Cool)

anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C blue no. 1, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum *may contain this ingredient

(Mucinex Nightshift Sinus Clear & Cool)

ammonium glycyrrhizate, anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C blue no. 1, flavors, glycerin (soy), propylene glycol, sodium benzoate, sorbitol, sucralose, water, xanthan gum

1-866-MUCINEX (1-866-682-4639)

Keep out of reach of children

Acetaminophen 650mg	Pain reliever
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10mg	

Acetaminophen 650mg......Pain reliever/fever reducer

Dextromethorphan HBr 20 mg.....Cough suppressant Triprolidine HCl 2.5 mg. Antihistamine



sucralose, water, xanthan gum Do not take Mudinex Sinus-Max Severe Congestion & Pain Clear & Cool and Mucinex Nightshift Sinus Clear & Cool at the same

time. Always wait at least 4 hours before taking another

dose of Mucinex liquid. TAKE

ONLY AS DIRECTED.





Drug Facts

Active ingredients (in each 20 mL) Purposes Mucinex Sinus-Max

SEVERE CONGESTION & PAIN CLEAR & COOL

Acetaminophen 650 mg	Pain reliever
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Drug Facts (continued)

Active ingredients (in each 20 mL) Purposes Mucinex Nightshift Sinus Clear & Cool

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant
Triprolidine HCl 2.5 mg	Antihistamine 🔻

Drug Facts (continued)

Uses Mucinex Sinus-Max SEVERE CONGESTION & PAIN CLEAR & COOL

- temporarily relieves:
- nasal congestion
- headache minor aches and pains
- sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Mucinex Nightshift Sinus Clear & Cool

- temporarily relieves these common cold and flu symptoms:

- nasal congestion
 minor aches and pains
- sore throat headache
- runny nose
- sinus congestion and pressure ■ sneezing ■ itching of the nose or throat
- itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning (Nightshift Sinus Clear & Cool only): If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
 heart disease
 diabetes
- high blood pressure
- thyroid disease
- glaucoma (Nightshift Sinus Clear & Cool only)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis (Nightshift Sinus Clear & Cool only)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Drug Facts (continued)

cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (Nightshift Sinus Clear & Cool only)

When using this product do not use more than directed

- excitability may occur, especially in children (Nightshift Sinus) Clear & Cool only)
- marked drowsiness may occur (Nightshift Sinus Clear & Cool only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (Nightshift Sinus Clear & Cool only)
- avoid alcoholic drinks (Nightshift Sinus Clear & Cool only)
- use caution when driving a motor vehicle or operating machinery (Nightshift Sinus Clear & Cool only)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

MUCINEX SINUS-MAX SEVERE CONGESTION & PAIN CLEAR & COOL

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

MUCINEX NIGHTSHIFT SINUS CLEAR & COOL

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

MUCINEX-SINUS MAX SEVERE CONGESTION AND PAIN CLEAR AND COOL AND MUCINEX NIGHTSHIFT SINUS CLEAR AND COOL

acetaminophen, dextromethorphan hydrobromide, quaifenesin, phenylephrine hydrochloride, triprolidine hydrochloride kit

Product Information

Product Type HUMAN OTC DRUG	Item Code (Source)	NDC:63824-116
-----------------------------	--------------------	---------------

II	Packaging					
	# Item Code	Package Description	Marketing Start Date	Marketing End Date		
	1 NDC:63824-116-66	1 in 1 CARTON	01/01/2019	01/29/2024		
	1	1 in 1 KIT				

Quant	Quantity of Parts		
Part #	Package Quantity	Total Product Quantity	
Part 1	1 BOTTLE	180 mL	
Part 2	1 BOTTLE	180 mL	

Part 1 of 2

MAXIMUM STRENGTH MUCINEX SINUS-MAX SEVERE CONGESTION AND PAIN CLEAR AND COOL

acetaminophen, guaifenesin, and phenylephrine hydrochloride solution

Product Information	
Item Code (Source)	NDC:63824-267
Route of Administration	ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg in 20 mL	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL	

Inactive Ingredients		
Ingredient Name	Strength	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)		
PROPYL GALLATE (UNII: 8D4SNN7V92)		
XANTHAN GUM (UNII: TTV12P4NEE)		
WATER (UNII: 059QF0KO0R)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SORBITOL (UNII: 506T60A25R)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		

D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics			
Color	green	Score	
Shape		Size	
Flavor	MENTHOL	Imprint Code	
Contains			

ı	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
		NDC:63824- 267-66	180 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	01/01/2019		

Part 2 of 2

MUCINEX NIGHTSHIFT SINUS CLEAR AND COOL

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, and triprolidine hydrochloride solution

Product Information				
Item Code (Source)	NDC:63824-270			
Route of Administration	ORAL			

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL		
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII: 2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	2.5 mg in 20 mL		
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg in 20 mL		
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL		

Inactive Ingredients				
Ingredient Name	Strength			
SUCRALOSE (UNII: 96K6UQ3ZD4)				
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
EDETATE DISODIUM (UNII: 7FLD91C86K)				
GLYCERIN (UNII: PDC6A3C0OX)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
SORBITOL (UNII: 506T60A25R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
WATER (UNII: 059QF0KO0R)				
XANTHAN GUM (UNII: TTV12P4NEE)				

Product Characteristics					
Color	green	Score			
Shape		Size			
Flavor	MENTHOL	Imprint Code			
Contains					

Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
		180 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)			

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC monograph final	part341	01/01/2019			

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC monograph final	part341	01/01/2019	01/29/2024		

Labeler - RB Health (US) LLC (081049410)

Revised: 8/2023 RB Health (US) LLC